


REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE	3. REPORT TYPE AND DATES COVERED FINAL REPORT 01 Sep 95 - 30 Nov 96	
4. TITLE AND SUBTITLE (DURIP-95) Research Instrumentation for Investigations in Atomic, Molecular and Optical Physics			5. FUNDING NUMBERS 61103D 3484/US	
6. AUTHOR(S) Dr Kenneth A. Hardy				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Physics Florida International University Miami, FL 33199			8. PERFORMING ORGANIZATION AFOSR-TR-97 0173	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR/NE 110 Duncan Avenue Suite B115 Bolling AFB DC 20332-8050			AGENCY REPORT NUMBER F49620-95-1-0515	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) A ring Ti/dye laser was purchased from Coherent Laser, model 899-21. The laser has been used in experiments involving metastable atom beam cooling. During this work the investigators developed a method of measuring laser line width that can be directly related to the fundamental quantities time and distance. Two spectra taken with the ring laser are included with this report. 19970602 040				
14. SUBJECT TERMS			15. NUMBER OF PAGES	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT	

DISCLAIMER NOTICE



**THIS DOCUMENT IS BEST
QUALITY AVAILABLE. THE
COPY FURNISHED TO DTIC
CONTAINED A SIGNIFICANT
NUMBER OF PAGES WHICH DO
NOT REPRODUCE LEGIBLY.**

FINAL TECHNICAL REPORT
RESEARCH INSTRUMENTATION FOR INVESTIGATIONS IN ATOMIC,
MOLECULAR AND OPTICAL PHYSICS

95-1-0515
USAFOSR GRANT # F49620-~~95-0515~~

KENNETH A. HARDY
FLORIDA INTERNATIONAL UNIVERSITY PHYSICS DEPARTMENT

A ring Ti/dye laser was purchased from Coherent Laser, model 899-21. The laser was delivered approximately February 1, 1996. The laser was not accepted at this time, however, due to difficulties with the specified line width. The investigators worked with the designers of the laser system to resolve the problems and the manufacturer agreed to redesign two circuit boards in the control system. This resulted in the laser system meeting the required specifications. The redesigned boards were delivered in July 1996.

The laser has been used in experiments involving metastable atom beam cooling. During this work the investigators developed a method of measuring laser line width that can be directly related to the fundamental quantities time and distance. We believe this method may have broad future application. The work is in preparation for publication.

Two spectra taken with the ring laser are included with this report. Figure 1 shows the result of laser cooling a beam of metastable argon atoms from a velocity of ≈ 750 m/sec to a velocity of ≈ 240 m/sec. The velocity width of the cooled atom beam has been reduced to 6%. Figure 2 shows laser cooling with the laser detuned 200 MHz. This gives a final velocity of 162 m/sec, which is approximately the low velocity limit of our apparatus in its current configuration. Figure two also illustrates the result of the laser cooling as it shows time spectra with the laser on and off. The two spectra shown were taken with a time channel width of 40 μ secs and a path length of 1.8 m. The final velocities are calculated knowing that the atoms travel .31 m with their initial velocity, a constant deceleration for .77 m and a final path length of .73 m, at constant final velocity.

Currently a magnetio-optical trap is being added to the apparatus to confine the cool atoms after which scattering experiments will commence with the goal of measuring the "S" wave scattering length of various metastable atom partners.




Figure 2 Hollow circles - TOF Distribution of the metastable atoms without the laser.
 Filled circles - TOF Distribution when the laser is detuned 200 MHz Red from resonance, and TMF with the currents 1.1, .7, .67, .61, .58, .45, .3 is used. Initial magnetic field 420 Gauss.
 Total TOF=5.94ms




Figure 1 TOF Distribution of metastable atoms when the laser is detuned 300 MHz Red from resonance and a TMF with $E_0 = 170$ Gauss is used.

TOF=5.64 ms